



Galaxea fascicularis
Octopus coral, fluorescence grass coral

Amir, H.

Citation: Amir, H., 2022. *Galaxea fascicularis*, Octopus coral, fluorescence grass coral. The Maldives Red List of Threatened Species.


Maldives National Red List Assessment: *Galaxea fascicularis*

A. Background Information

1. Assessment Information:

Assessor Name(s)	Hana Amir
Date of Assessment	10 February 20222
Contributors	Phillipa Roe, Simone Montano, Ilham Mohamed
Institutional contributors	Ministry of Environment, Climate Change and Technology; Maldives Marine Research Institute; IUCN Project REGENERATE; Maldives Allen Coral Atlas; Noo Raajje; Six Senses Laamu
Facilitators	James Tallant
Reviewers	Janice Chanson, Maldives Marine Research Institute

2. Taxonomic Information:

Scientific Name	<i>Galaxea fascicularis</i> (Linnaeus, 1767)	
Common Name (English)	Octopus coral, fluorescence grass coral	
Common Name (Dhivehi)	No species name Generic name: Ufuli gaa 	
Taxonomy	Order	Scleractinia
	Sub-order	-
	Family	Euphylliidae
Taxonomic Notes	Originally <i>Madrepora fascicularis</i> (Linnaeus, 1767) Formerly of family Oculinidae	

3. Geographic Range:

3.1 Summary of Global Distribution


The species is found in all parts of the Indian Ocean stretching all the way to Australia, Japan, China, and the Pacific. It is also found in the Red Sea and the Gulf of Aden (Hoeksema et al., 2014).

3.2 Countries of Occurrence

American Samoa, Australia, Bahrain, British Indian Ocean Territory, Cambodia, Comoros, Cook Islands, Djibouti, Egypt, Eritrea, Fiji, French Polynesia, Guam, India, Indonesia, Iran, Islamic Republic of Iraq, Israel, Japan, Jordan, Kenya, Kiribati, Kuwait, Madagascar, Malaysia, Maldives, Marshall Islands, Mauritius, Mayotte, Micronesia, Federated States of Mozambique, Myanmar, Nauru, New Caledonia, Niue, Northern Mariana Islands, Oman, Palau, Papua New Guinea, Philippines, Qatar, Reunion, Samoa, Saudi Arabia, Seychelles, Singapore, Solomon Islands, Somalia, Sri Lanka, Sudan, Taiwan, Tanzania, United Republic Thailand, Tokelau, Tonga, Tuvalu, United Arab Emirates, United States Minor Outlying Islands, Vanuatu, Vietnam, Wallis and Futuna, Yemen (Hoeksema et al. 2014)

3.3 National distribution

Exact species distribution is unknown. There are confirmed records of the species (Pichon and Benzoni 2007) and the genus has been recorded from most atolls within the Maldives. Based on observations, confirmed genus data, available data, and species characteristics, it is estimated that the genus is widely distributed from the North to the South of the Maldives. Hence, depending on the depth and habitat, it is possible that species is also widely distributed across all the atolls.



4. Population:

4.1 Summary

There is no species-specific population information available. Therefore, population size, trends and distribution are estimated from generic information.

4.2 Population Size

Global population is generalized as “common” (Devantier, Turak and Szava-Kovats 2020). National species population size is unknown.

A ten-year generic mean, the mean percent cover of the genus over a ten year period of time, was calculated for each genus. This was used as a proxy for the potential species population size.

The ten-year generic mean derived from available data is $0.54 \pm 0.70\%$ (\pm SE). Recent estimates indicate cover is between >0.00 and $\sim 5.00\%$ (Noo Raajje 2021). Therefore, based on generic estimates the species population may vary between $\sim >0.0$ to $\sim 5\%$ cover assuming differing levels of dominance in a community.

4.3 Population Trend

Global and national past population trends are not known. However, although the species is thought to be relatively resistant to bleaching, local records indicate that the species is highly impacted by thermal stress anomalies and is vulnerable to bleaching. Future population trends are therefore likely to be decreasing.

4.4 Generation length

Estimates for coral generation lengths are based on Carpenter *et al.* 2008, as follows. The coral colony is considered to be the mature individual, as it typically lives, is injured, or dies as a unit. The average age of natural survival of a coral colony was defined as the average age of a mature individual, or one generation length. Based on available knowledge of coral species' biology and life history, this was determined to be 10 years. Therefore, population reductions are estimated over 30 years, representing three generation lengths.

4.5 Continuing Decline in Number of Mature Individuals?

Unknown

4.6 Extreme Fluctuations?

No

4.7 Severely Fragmented?

No. Population assumed to be cohesive based on generic distribution and mode of reproduction, although degree of connectivity (genetic or otherwise) within and between atolls is unknown.

5. Habitat and Ecology:

5.1 Summary

The species can be found on mid to low slopes on reef habitats between depths of 1-15 m. It is generally found in locations that are protected from exposure and wave action (Veron *et al.*, 2022).

5.2 Systems (terrestrial / freshwater / marine)

Marine

5.3 Continuing Decline in Area, Extent, and/or Quality of Habitat?

Yes. There is continuous loss of area, extent and habitat quality stemming from coastal development projects and increase in frequency and magnitude of climate change impacts including extreme temperature fluctuations.

5.4 A Migratory Species?

No

6. Use and Trade

6.1 Is the species used or traded?

Yes

6.2 Summary

The species is collected for aquarium use. This harvesting may represent a threat to the species.

7. Threats:

7.1 Summary

Species specific threats are unknown. However, the genus is known to be vulnerable to sedimentation with low recovery (Junjie et al., 2014) and coral disease (Ammar et al., 2011). Its susceptibility to bleaching is low (Baird et al. 2009) though recovery from bleaching variable (Li et al., 2012). As the species is found in shallow habitats (<15m), it is expected that it is vulnerable to both coastal development and climate change impacts. Moreover, its habitat is located close to human habitation, either island communities or resorts, and therefore they are subject to stressors in the form of untreated sewage (where infrastructure is lacking), agricultural runoff and/or domestic waste.

The projected date of onset of Annual Severe Bleaching (ASB) is the date after which the capacity of coral reefs to recover from repeated bleaching events is expected to significantly reduce - resulting in fundamental, permanent population changes (UNEP 2017). The calculation of ASB by UNEP (2020) is restricted to a 30 m depth, due to the dataset used for the spatial analysis.

We calculated the mean year of ASB onset, assuming 0° coral adaptation (UNEP 2020) to climate change, for areas occupied by the genus. A 0° adaptation (i.e. no adaptation) to climate change was applied as a precautionary approach:

- as the capacity of coral to adapt to climate change is not yet well understood
- as there are other climate change impacts that may compound population decline and affect population refuges, and
- to account for impacts from coastal development that could potentially exacerbate the effect of bleaching prompted by climate change.

Based on this analysis, *G. fascicularis* is expected to undergo ASB starting in 2041.

7.2 Major Threats to the Species (using IUCN Threat Classification Scheme)

1. Residential & commercial development:

1.1. Housing & urban areas

Timing: Ongoing

Stresses:

1. *Ecosystem/Community stresses*
 - 1.1. *Ecosystem conversion*
 - 1.2. *Ecosystem degradation*
 - 1.3. *Indirect ecosystem effects*
2. *Species stresses*
 - 2.2. *Species disturbance*
 - 2.3. *Indirect species effects*
 - 2.3.7. *Reduced reproductive success*

1.2. Commercial & industrial areas

Timing: Ongoing

Stresses:

1. *Ecosystem/Community stresses*
 - 1.1. *Ecosystem conversion*
 - 1.2. *Ecosystem degradation*
 - 1.3. *Indirect ecosystem effects*
2. *Species stresses*
 - 2.2. *Species disturbance*
 - 2.3. *Indirect species effects*
 - 2.3.7. *Reduced reproductive success*

1.3. Tourism & recreational areas

Timing: Ongoing

Stresses:

1. Ecosystem/Community stresses

1.1. Ecosystem conversion

1.2. Ecosystem degradation

1.3. Indirect ecosystem effects

2. Species stresses

2.2. Species disturbance

2.3. Indirect species effects

2.3.7. Reduced reproductive success

5. Biological resource use:

5.4. Fishing & harvesting aquatic resource

5.4.3. Unintentional effects: subsistence/small scale

Timing: Ongoing

Stresses:

1. Ecosystem/Community stresses

1.2. Ecosystem degradation

7. Natural system modifications:

7.3. Other ecosystem modifications

Timing: Ongoing

Stresses:

1. Ecosystem/Community stresses

1.1. Ecosystem conversion

1.2. Ecosystem degradation

9. Pollution:

9.1. Domestic & urban wastewater

9.1.1. Sewage

Timing: Ongoing

Stresses:

1. Ecosystem/Community stresses

1.2. Ecosystem degradation

9.1.2. Run-off

Timing: Ongoing

Stresses:

1. Ecosystem/Community stresses

1.2. Ecosystem degradation

9.4. Garbage and solid waste

Timing: Ongoing

Stresses:

1. Ecosystem/Community stresses

1.2. Ecosystem degradation

11. Climate change & severe weather:

11.1. Habitat shifting & alteration

Timing: Ongoing

Stresses:

1. Ecosystem/Community stresses

1.2. Ecosystem degradation

2. Species stresses

2.1. Species mortality

2.2. Species disturbance

2.3. Indirect species effects

2.3.2. Competition

2.3.7. Reduced reproductive success

11.4. Storms & flooding

Timing: Ongoing

Stresses:

7.3 Species vulnerability modifiers

Coral abundance and distribution data is only available at the genus level. Species characteristics were used to evaluate how susceptible each species is to treats. This data was gathered from the IUCN SSC Coral Working Group's Global Traits Database, and available literature. Each species was assigned a vulnerability score for each trait (3 = high vulnerability; 2 = medium vulnerability; 1 = low vulnerability). This species was scored as follows.

Trait	Vulnerability score
Susceptibility to bleaching	1
Resistance to bleaching	1
Recovery from bleaching or disease	2
Resistance to disease	2
Susceptibility to predation	2
Recovery from sedimentation	2
Resistance to sedimentation	1
Susceptibility to sedimentation	3
<i>Average</i>	2

8. Conservation and Research:

8.1 Summary

The species is listed as "Near Threatened" on the IUCN Red List of Threatened Species and is included within CITES Appendix II. Collection, killing and export of live and dead scleractinian corals is illegal under Maldivian fisheries law. Distribution of the species falls under multiple national MPAs though not all the MPAs are properly managed. Species distribution and population information would be required to better manage and protect the species especially as it is a popular target for propagation products and for aquaria. Further species-specific research for threats would support development and implementation of necessary legislation.

8.2 Conservation Actions/Research in Place

Globally, as a coral the species is included within CITES Appendix II. It is listed as "Near Threatened" within the IUCN Red List of Threatened Species.

Nationally, collection, killing, and export of both live and dead scleractinian corals (under phylum Cnidaria) are illegal in the Maldives under the Maldives Fisheries Law 2020/R-75. The distribution of the genus, and likely the species, overlaps multiple MPAs including Baa atoll biosphere reserve SAMPA.

8.3 Conservation Actions Needed

Despite laws and regulations protecting coral species within the country, implementation of such laws and regulations are a challenge. There is a need to create infrastructure and capacity that would enable relevant laws and regulations to be enforced whilst educating the general public that they exist. Educational and training needs extend to more educational opportunities in the marine field as well as skill training opportunities that can support research, further education, institutional development, awareness and communication efforts. Moreover, with the spatial distribution of

maldivian reefs and the corals that are found within them, there is an urgent need to foster intra-governmental as well as inter-agency and inter-organizational cooperation and partnerships to mediate limitations related to capacity and resources. Areas of high abundance of specific species along with areas of high coral cover and resilience need to be identified and managed if not fully protected to ensure sustainable use and longevity. Management includes mitigating impacts from coastal development and resolving conflict of multi-use resources. Further, with increasing interest in restoration of reefs, there is a need to properly manage species harvest for projects to ensure that wild stocks are not decimated and species are able to recover.

1. Land/water protection

- 1.1. Site/area protection
- 1.2. Resource & habitat protection

2. Land/water management

- 2.1. Site/area management
- 2.3. Habitat & natural process restoration

3. Species management

- 3.1. Species management
 - 3.1.1. Harvest management
- 3.2. Species recovery
- 3.4. *Ex situ* conservation
 - 3.4.1. Captive breeding/artificial propagation
 - 3.4.2. Genome resource bank

4. Education & awareness

- 4.1. Formal education
- 4.2. Training
- 4.3. Awareness & communications

5. Law & policy

- 5.3. Private sector standards & codes
- 5.4. Compliance & enforcement
 - 5.4.1. Scale unspecified

7. External capacity building

- 7.1. Institutional & civil society development
- 7.2. Alliance & partnership development
- 7.3. Conservation finance

8.4 Research Needed

1. Research

- 1.2. Population size, distribution and past trends
- 1.3. Life history and ecology
- 1.5. Threats
- 1.6. Conservation actions

3. Monitoring

- 3.1. Population trends
- 3.4. Habitat trends

B. Assessment

9. CRITERION A

Criterion A

Generation Length		10 years
3 Generations		30 years
Reduction in population size?		Yes
Start Date of Reduction		2022
End Date of Reduction		2052
Rate of Reduction (%)		30-40%
Meets Criteria Thresholds?	A1	-
	A2	-
	A3	Projected 30-40% reduction over the next three generations (VU A3)
	A4	-
Reduction Based on Which Sub-criteria?	a	-
	b	-
	c	Decline in habitat quality associated with impacts of climate change
	d	-
	e	Mortality due to repeated bleaching

Assessment Under Criterion A

Reasoning Our assessment of future population reduction is based on the projected date of onset of ASB. *G. fascicularis* is expected to experience ASB starting in 2041.

The species ranges between 1-15 m and is fully restricted to depths shallower than 30 m. Depth and depth range is considered in our assessment, as populations at depths shallower than 30 m are expected to experience greater temperature fluctuations and extreme temperatures (Riegl and Piller 2003), and therefore decline more quickly.

This genus was noted to be one that bleached intensely in the Maldives in 2016 (Ibrahim et al. 2017), suggesting that it is relatively vulnerable to thermal stress.

However, this species has also been noted to be relatively resilient to bleaching, with low susceptibility and high resistance based on literature review and scientific expertise. Therefore, while a decline is expected over 100% of the species' depth range, this reduction is likely not to be as severe as that of other coral species..

Therefore, this species is projected to undergo a reduction of 30-40% over the next three generations (2022-2052).

10. CRITERION B

Criterion B

AOO		-
EEO		-
Meets Criteria Thresholds?	B1	-
	B2	-
Severely Fragmented?		-
No. Locations		-
Threat used to calculate locations		-
Continuing Decline?	(i) EEO	-
	(ii) AOO	-
	(iii) Habitat	-
	(iv) Locations / Subpopulations	-
	(v) Mature Individuals	-
Extreme Fluctuations?	(i) EEO	-
	(ii) AOO	-
	(iii) Locations / Subpopulations	-
	(iv) Mature Individuals	-

Assessment Under Criterion B DD

Criterion C		
No. Mature Individuals		Unknown
Continuing Decline in Population Size?		-
Is Rate of Decline Known?		-
Generation Length		-
C1	Meets Thresholds for Rate of Decline?	-
	Rate of Decline (%)	-
	Time Period of Decline	-
C2	(a) (i) No. Mature Individuals in Each Subpopulation	-
	(a) (ii) % Mature Individuals in one Sub-population	-
	(b) Extreme Fluctuations in No. Mature Individuals?	-
Assessment Under Criterion C		DD

Criterion D		
No. Mature Individuals		Unknown
Meets Criteria Thresholds?		-
VUD2	AOO	-
	No. Locations	-
	Plausible Future Threat That Would Quickly Drive Taxon to CR or EX	-

	Plausible Future Threat That Would Quickly Drive Taxon to VU or EN	-
Assessment Under Criterion D		DD

Criterion E	
Has a Quantitative Analysis Been Conducted?	No
Type of Quantitative Analysis	-
Generation Length	-
Probability of Extinction within 100 Years	-
Probability of Extinction within 20 Years / 5 Generations (whichever is longer)	-
Probability of Extinction within 10 Years / 3 Generations (whichever is longer)	-
Assessment Under Criterion E	
DD	

Preliminary Assessment		VU A3ce
Regional Adjustment	Up-list, Down-list, or No Change?	No change
	Justification for Regional Adjustment	No likely interaction with individuals outside the region
Final Assessment		VU A3ce
Narrative Justification for Assessment:		
<p><i>Galaxea fascicularis</i> is likely distributed throughout all the atolls of the Maldives, with a depth range of 1-15 m. Coral species found at depths of less than 30 m are more exposed to extreme and fluctuating water temperatures associated with the impacts of climate change, which can lead to population reductions.</p> <p>Although the genus suffered in the most recent bleaching event in the Maldives, and it is vulnerable to disease and sedimentation, the species has been identified as being relatively resilient to bleaching (with high resistance and low susceptibility).</p> <p>While the current population size and past rate of decline are not known, it is anticipated that the population will undergo a significant decline in the future based on the impacts of climate change on its local range. Based on published climate model-based bleaching assessments published by UNEP (2017, 2020), it is estimated that ASB will commence in the year 2041 over this species' distribution, affecting 100% of its depth range. However, as this species is known to be relatively resilient to bleaching, we anticipate that this will constitute a population reduction of only 30-40%.</p> <p>Therefore, based on the best available data, we project a 30-40% reduction in population size within the next three generations (2022-52), prompted by degradation in habitat quality associated with the impacts of climate change. This meets the threshold for Vulnerable under Criterion A3c.</p>		

C. References

- Ammar, M. S., Obuid-Allah, A. H., & Al-Hammady, M. A. (2011). Corals differential susceptibilities to bleaching along the Red Sea Coast, Egypt. *Nusantara Bioscience*, 3(2).
- Baird, A. H., Bhagooli, R., Ralph, P. J., & Takahashi, S. (2009). Coral bleaching: the role of the host. *Trends in Ecology & Evolution*, 24(1), 16-20.
- Carpenter, K.E., Abrar, M., Aeby, G., Aronson, R.B., Banks, S., Bruckner, A., Chiriboga, A., Cortés, J., Delbeek, J.C., DeVantier, L., Edgar, G.J., Edwards, A.J., Fenner, D., Guzmán, H.M., Hoeksema, B.W., Hodgson, G., Johan, O., Licuanan, W.Y., Livingstone, S.R., Lovell, E.R., Moore, J.A., Obura, D.O., Ochavillo, D., Polidoro, B.A., Precht, W.F., Quibilan, M.C., Reboton, C., Richards, Z.T., Rogers, A.D., Sanciangco, J., Sheppard, A., Sheppard, C., Smith, J., Stuart, S., Turak, E., Veron, J.E.N., Wallace, C., Weil, E. and Wood, E. 2008. One-Third of Reef-Building Corals Face Elevated Extinction Risk from Climate Change and Local Impacts. *Science*. 25 July 2008: 560-563. DOI: 10.1126/science.1159196. Supporting online material: www.sciencemag.org/cgi/content/full.1159196.
- DeVantier, L., & Turak, E. (2017). Species richness and relative abundance of reef-building corals in the Indo-West Pacific. *Diversity*, 9(3), 25.
- DeVantier, L., Turak, E., & Szava-Kovats, R. (2020). Species Richness and Abundance of Reef-Building Corals in the Indo-West Pacific: The Local–Regional Relation Revisited. *Frontiers in Marine Science*, 7, 487.
- Hoeksema, B. W.; Cairns, S. (2022). World List of Scleractinia. *Galaxea fascicularis* (Linnaeus, 1767). Accessed through: World Register of Marine Species at: <https://www.marinespecies.org/aphia.php?p=taxdetails&id=207366> on 2022-02-05
- Hoeksema, B.W., Rogers, A. & Quibilan, M.C. 2014. *Galaxea fascicularis*. *The IUCN Red List of Threatened Species* 2014: e.T133662A54303932. <https://dx.doi.org/10.2305/IUCN.UK.2014-1.RLTS.T133662A54303932.en>. Accessed on 04 March 2022.
- Ibrahim, N., Mohamed, M., Basheer, A., Haleem, I., Nistharan, F., Naeem, R., Abdulla, A., & Grimsditch, G. (2017). *Status of Coral Bleaching in the Maldives in 2016*.
- Junjie, R. K., Browne, N. K., Erfemeijer, P. L., & Todd, P. A. (2014). Impacts of sediments on coral energetics: partitioning the effects of turbidity and settling particles. *PLoS One*, 9(9), e107195.
- Li, X., Liu, S., Huang, H., Huang, L., Jing, Z., & Zhang, C. (2012). Coral bleaching caused by an abnormal water temperature rise at Luhuitou fringing reef, Sanya Bay, China. *Aquatic Ecosystem Health & Management*, 15(2), 227-233.
- NooRaajje (2021). Maldives Coral Reef Assessment. *Waïtt Institute*. 73 pp.
- Pichon, M., & Benzoni, F. (2007). Taxonomic re-appraisal of zooxanthellate Scleractinian Corals in the Maldivian Archipelago. *Zootaxa*, 33(1441), 21–33. <https://doi.org/10.11646/zootaxa.1441.1.2>
- Rasheed, S., Warder, S. C., Plancherel, Y., & Piggott, M. D. (2021). An Improved Gridded Bathymetric Data Set and Tidal Model for the Maldives Archipelago. *Earth and Space Science*. <https://doi.org/10.1029/2020EA001207>
- Riegl, B. and Piller, W.E. 2003. Possible refugia for reefs in times of environmental stress. *International Journal of Earth Sciences* 92(4):520-531.

UNEP 2017. Coral Bleaching Futures - Downscaled projections of bleaching conditions for the world's coral reefs, implications of climate policy and management responses. United Nations Environment Programme, Nairobi, Kenya

UNEP 2020. Projections of future coral bleaching conditions using IPCC CMIP6 models: climate policy implications, management applications, and Regional Seas summaries. United Nations Environment Programme, Nairobi, Kenya

Veron J.E.N., Stafford-Smith M.G., Turak E. and DeVantier L.M. (2022). Corals of the World. Accessed 9 February 2022, Version 0.01 (Beta). [http://coralsoftheworld.org/v0.01\(Beta\)](http://coralsoftheworld.org/v0.01(Beta)). (To go to the current version access: <http://coralsoftheworld.org>)