



Isopora palifera

Montano, S.

Citation: Montano, S., 2022. *Isopora palifera*. The Maldives Red List of Threatened Species.

Maldives National Red List Assessment: *Isopora palifera*

A. Background Information

1. Assessment Information:

Assessor Name	Simone Montano
Date of Assessment	17/02/2022
Contributors	Hana Amir, Phillipa Roe, Ilham Mohamed
Institutional contributors	Ministry of Environment, Climate Change and Technology; Maldives Marine Research Institute; IUCN Project REGENERATE; Maldives Allen Coral Atlas; Noo Raajje; Six Senses Laamu, MarHE center
Facilitators	James Tallant
Reviewers	Janice Chanson, Maldives Marine Research Institute

2. Taxonomic Information:

Scientific Name	<i>Isopora palifera</i> (Lamarck, 1816)	
Common Name (English)	-	
Common Name (Dhivehi)	-	
Taxonomy	Order	Scleractinia
	Sub-order	-
	Family	Acroporidae
Taxonomic Notes	-	

3. Geographic Range:

3.1 Summary of Global Distribution

Distinct populations can be found in the Indo-Pacific

3.2 Countries of Occurrence

American Samoa; Australia; British Indian Ocean Territory; Cambodia; Christmas Island; Cocos (Keeling) Islands; Comoros; Djibouti; Eritrea; Fiji; Guam; India; Indonesia; Japan; Kenya; Kiribati; Madagascar; Malaysia; Maldives; Marshall Islands; Mauritius; Mayotte; Micronesia, Federated States of ; Mozambique; Myanmar; Nauru; New Caledonia; Northern Mariana Islands; Palau; Papua New Guinea; Philippines; Réunion; Samoa; Seychelles; Singapore; Solomon Islands; Somalia; South Africa; Sri Lanka; Taiwan, Province of China; Tanzania, United Republic of; Thailand; Tuvalu; Vanuatu; Viet Nam; Wallis and Futuna; Yemen

3.3 National Distribution

Although exact species distribution is unknown, presence records for the species are available in literature and the genus has been observed in the country.

Data confirmed occurrences of the genus are not available from every atoll. Based on known observations, available data, species characteristics it is estimated that there is possibility for the genus and the species to occur on shallow reef systems in the North, Central and South of the Maldives. Depending on depth, the species may not be found.

4. Population:

4.1 Summary

Species specific population information is not available. Population estimates and distribution are derived based on best available generic information.

4.2 Population Size

Global population is generalised as Common. National species population size is unknown.

4.3 Population Trend

Past population trends are unknown. The future population trend is likely to be decreasing.

4.4 Generation length

Estimates for coral generation lengths are based on Carpenter *et al.* 2008, as follows. The coral colony is considered to be the mature individual, as it typically lives, is injured, or dies as a unit. The average age of natural survival of a coral colony was defined as the average age of a mature individual, or one generation length. Based on available knowledge of coral species' biology and life history, this was determined to be 10 years. Therefore, population reductions are estimated over 30 years, representing three generation lengths.

4.5 Continuing Decline in Number of Mature Individuals?

Unknown

4.6 Extreme Fluctuations?

No

4.7 Severely Fragmented?

No. Population assumed to be cohesive based on generic distribution although degree of connectivity between atolls is unknown.

5. Habitat and Ecology:

5.1 Summary

Frequently found between 2 m and 20 m depths, this species occurs in all shallow, tropical reef environments. It is found subtidally on deeper parts of reef flat and reef edge, reef slopes and walls, and submerged reefs (Wallace 1999).

5.2 Systems (terrestrial / freshwater / marine)

Marine

5.3 Continuing Decline in Area, Extent, and/or Quality of Habitat?

Yes. There is continuous loss of area, extent and habitat quality stemming from coastal development projects and increase in frequency and magnitude of climate change impacts including extreme temperature fluctuations.

5.4 A Migratory Species?

The species is not migratory

6. Use and Trade:

6.1 Is the species used or traded?

No

6.2 Summary

-

7. Threats:

7.1 Summary

The understanding of threats to this species are predominantly based on genus-level threats. The genus is known to be vulnerable to sedimentation, bleaching, predation by corallivores (e.g., COTS) and disease (BrB). The species is affected by overall declines in ecosystem and habitat quality. Therefore, as the species is found in shallow reef habitats (<15 m), it is expected that the species is particularly vulnerable to the impacts of coastal development projects and of climate change. Finally, this is a brooder and therefore has a smaller sexual reproductive output and limited dispersal capacity.

The projected date of onset of Annual Severe Bleaching (ASB) is the date after which the capacity of coral reefs to recover from repeated bleaching events is expected to significantly reduce - resulting in fundamental, permanent population changes (UNEP 2017). The calculation of ASB by UNEP (2020) is restricted to a 30 m depth, due to the dataset used for the spatial analysis.

We calculated the mean year of ASB onset, assuming 0° coral adaptation (UNEP 2020) to climate change, for areas occupied by the genus. A 0° adaptation (i.e. no adaptation) to climate change was applied as a precautionary approach:

- as the capacity of coral to adapt to climate change is not yet well understood
- as there are other climate change impacts that may compound population decline and affect population refuges, and
- to account for impacts from coastal development that could potentially exacerbate the effect of bleaching prompted by climate change.

Based on this analysis, *I. palifera* is expected to undergo ASB starting in 2041.

7.2 Major Threats to the Species (using IUCN Threat Classification Scheme)

Residential & commercial development:

- 1.1. Housing & urban areas
- 1.2. Commercial & industrial areas
- 1.3. Tourism & recreational areas

Natural system modifications:

- 7.3. Other ecosystem modifications

Invasive & other problematic species, genes & disease:

- 8.2. Problematic native species/diseases
 - 8.2.1. Unspecified species/disease

Climate change & severe weather:

- 11.1. Habitat shifting & alteration
- 11.2. Temperature extremes
- 11.4. Storms & flooding

7.3 Species vulnerability modifiers

Coral abundance and distribution data is only available at the genus level. Species characteristics were used to evaluate how susceptible each species is to treats. This data was gathered from the IUCN SSC Coral Working Group's Global Traits Database, and available literature. Each species was assigned a vulnerability score for each trait (3 = high vulnerability; 2 = medium vulnerability; 1 = low vulnerability). This species was scored as follows.

Trait	Vulnerability score
Susceptibility to bleaching	3
Resistance to bleaching	3
Recovery from bleaching or disease	3
Resistance to disease	3
Susceptibility to predation	3
Recovery from sedimentation	2
Resistance to sedimentation	2
Susceptibility to sedimentation	2
<i>Average</i>	3

8. Conservation and Research:

8.1 Summary

The species is listed as "Near Threatened" on the IUCN Red List of Threatened Species and is included within the CITES appendix II. Collection, killing and export of live and dead scleractinian corals is illegal under Maldivian fisheries law. Distribution of the species falls under multiple national MPAs though not all the MPAs are properly managed. Species distribution and population information would be required to better manage and protect the species and further species-specific research for threats would support development and implementation of necessary legislature.

8.2 Conservation Actions/Research in Place

Globally, the species is listed within CITES Appendix II and listed as Near Threatened within the IUCN Red List of Threatened Species.

8.3 Conservation Actions Needed

Recommended measures for conserving this species include research in abundance and trends, ecology and habitat status, threats and resilience to threats. Establishment and management of new protected areas; expansion of protected areas; recovery management; and disease, pathogen and parasite management are strongly suggested. Restoration action may include artificial asexual and sexual propagation. These strategies will become important for conserving coral biodiversity.

1. Land/water protection
 - 1.1. Site/area protection
 - 1.2. Resource & habitat protection
2. Land/water management
 - 2.1. Site/area management
 - 2.3. Habitat & natural process restoration
4. Education & awareness
 - 4.1. Formal education
 - 4.2. Training
 - 4.3. Awareness & communications

8.4 Research Needed

1. Research
 - 1.2. Population size, distribution and past trends
 - 1.3. Life history and ecology
 - 1.5. Threats
 - 1.6. Conservation actions
3. Monitoring
 - 3.1. Population trends
 - 3.4. Habitat trends

B. Assessment

9. CRITERION A

Criterion A

Generation Length	10 years
3 Generations	30 years
Reduction in population size?	Yes
Start Date of Reduction	2022
End Date of Reduction	2052
Rate of Reduction (%)	80-90%
Meets Criteria Thresholds?	A1
	A2
	A3 Projected reduction in population size of 80-90% over the next three generations (CR A3).
	A4
Reduction Based on Which Sub-criteria?	a
	b
	c Decline in habitat quality associated with impacts of climate change.
	d
	e Mortality due to repeated bleaching.

Assessment Under Criterion A CR A3ce

Reasoning

Our assessment of future population reduction is based on the projected date of onset of ASB. *I. palifera* is expected to undergo ASB starting in 2041.

The species ranges between 2 m and 20 m. Depth and depth range is considered in our assessment, as populations at depths shallower than 30 m are expected to

experience greater temperature fluctuations and extreme temperatures (Riegl and Piller 2003), and therefore decline more quickly.

In addition, the species is considered to have low resistance to bleaching, with high susceptibility to the same threat based on literature review and scientific expertise. Similarly, it is susceptible to diseases and predation.

This species is projected to undergo a population reduction of 80-90% over the next three generations (2022-2052).

10. CRITERION B

Criterion B

AOO		-
EOO		-
Meets Criteria Thresholds?	B1	-
	B2	-
Severely Fragmented?		-
No. Locations		-
Threat used to calculate locations		-
Continuing Decline?	(i) EOO	-
	(ii) AOO	-
	(iii) Habitat	-
	(iv) Locations / Subpopulations	-
	(v) Mature Individuals	-
Extreme Fluctuations?	(i) EOO	-
	(ii) AOO	-
	(iii) Locations / Subpopulations	-
	(iv) Mature Individuals	-

Assessment Under Criterion B DD

11. CRITERION C

Criterion C

No. Mature Individuals		Unknown
Continuing Decline in Population Size?		-
Is Rate of Decline Known?		-
Generation Length		-
C1	Meets Thresholds for Rate of Decline?	-
	Rate of Decline (%)	-
	Time Period of Decline	-
C2	(a) (i) No. Mature Individuals in Each Subpopulation	-
	(a) (ii) % Mature Individuals in one Sub-population	-
	(b) Extreme Fluctuations in No. Mature Individuals?	-

Assessment Under Criterion C DD

12. CRITERION D

Criterion D

No. Mature Individuals		Unknown
Meets Criteria Thresholds?		-
VUD2	AOO	-
	No. Locations	-
	Plausible Future Threat That Would Quickly Drive Taxon to CR or EX	-

Plausible Future Threat That Would Quickly Drive Taxon to VU or EN -

Assessment Under Criterion D DD

13. CRITERION E

Criterion E

Has a Quantitative Analysis Been Conducted?	No
Type of Quantitative Analysis	-
Generation Length	-
Probability of Extinction within 100 Years	-
Probability of Extinction within 20 Years / 5 Generations (whichever is longer)	-
Probability of Extinction within 10 Years / 3 Generations (whichever is longer)	-

Assessment Under Criterion E DD

14. Preliminary and Final Assessment

Preliminary Assessment

CR A3ce

Regional Adjustment	Up-list, Down-list, or No Change?	No change
	Justification for Regional Adjustment	No likely interaction with individuals outside the region

Final Assessment CR A3ce

Narrative Justification for Assessment:

Isopora palifera is likely distributed throughout all the atolls of the Maldives, with a depth range of 2-30 m. Coral species found at depths of less than 30 m are more exposed to extreme and fluctuating water temperatures associated with the impacts of climate change, which can lead to population reductions.

Members of this genus are known to have a low resistance and low tolerance to bleaching and disease, and be slow to recover. A key threat to this species is extensive reduction in coral reef habitat, through a combination of processes.

Although its population size is not known, we anticipate that this species will undergo a severe decline in the future based on the impacts of climate change on its local range. Based on published climate model-based bleaching assessments published by UNEP (2017, 2020), Annual Severe Bleaching (ASB) is likely to commence in 2041 over this species' distribution, affecting 100% of its depth range. ASB reduces corals' ability to recover from repeated bleaching events, resulting in fundamental, permanent population changes (UNEP 2017). ASB is expected to constitute at least a 80-90% reduction in population size.

Therefore, we project a 80-90% population reduction over the next three generations (2022-2052), resulting from degradation in habitat quality associated with the impacts of climate change. This meets the threshold for CR A3c.

C. References

Carpenter, K.E., Abrar, M., Aeby, G., Aronson, R.B., Banks, S., Bruckner, A., Chiriboga, A., Cortés, J., Delbeek, J.C., DeVantier, L., Edgar, G.J., Edwards, A.J., Fenner, D., Guzmán, H.M., Hoeksema, B.W., Hodgson, G., Johan, O., Licuanan, W.Y., Livingstone, S.R., Lovell, E.R., Moore, J.A., Obura, D.O., Ochavillo, D., Polidoro, B.A., Precht, W.F., Quibilan, M.C., Reboton, C., Richards, Z.T., Rogers, A.D., Sanciangco, J., Sheppard, A., Sheppard, C., Smith, J., Stuart, S., Turak, E., Veron, J.E.N., Wallace, C., Weil, E. and Wood, E. 2008. One-Third of Reef-Building Corals Face Elevated Extinction Risk from Climate Change and Local Impacts. *Science*. 25 July 2008: 560-563. DOI: 10.1126/science.1159196. Supporting online material: www.sciencemag.org/cgi/content/full.1159196.

Montano, S., Strona, G., Seveso, D., Maggioni, D., & Galli, P. (2015). Widespread occurrence of coral diseases in the central Maldives. *Marine and Freshwater Research*, 67(8), 1253-1262.

Montano, S., Giorgi, A., Monti, M., Seveso, D., & Galli, P. (2016). Spatial variability in distribution and prevalence of skeletal eroding band and brown band disease in Faafu Atoll, Maldives. *Biodiversity and Conservation*, 25(9), 1625-1636.

Preston, S., & Richards, Z. (2021). Biological consequences of an outbreak of growth anomalies on *Isopora palifera* at the Cocos (Keeling) Islands. *Coral Reefs*, 40(1), 97-109.

Sheppard, C., Sheppard, A., & Fenner, D. (2020). Coral mass mortalities in the Chagos Archipelago over 40 years: regional species and assemblage extinctions and indications of positive feedbacks. *Marine pollution bulletin*, 154, 111075.

UNEP 2017. *Coral Bleaching Futures - Downscaled projections of bleaching conditions for the world's coral reefs, implications of climate policy and management responses*. United Nations Environment Programme, Nairobi, Kenya

UNEP 2020. *Projections of future coral bleaching conditions using IPCC CMIP6 models: climate policy implications, management applications, and Regional Seas summaries*. United Nations Environment Programme, Nairobi, Kenya

Wallace, C.C. 1999. *Staghorn Corals of the World: a revision of the coral genus Acropora*. CSIRO, Collingwood.